Parkes Weber syndrome

Parkes Weber syndrome is a disorder of the vascular system, which is the body's complex network of blood vessels. The vascular system consists of arteries, which carry oxygen-rich blood from the heart to the body's various organs and tissues; veins, which carry blood back to the heart; and capillaries, which are tiny blood vessels that connect arteries and veins.

Parkes Weber syndrome is characterized by vascular abnormalities known as capillary malformations and arteriovenous fistulas (AVFs), which are present from birth. The capillary malformations increase blood flow near the surface of the skin. They usually look like large, flat, pink stains on the skin, and because of their color are sometimes called "port-wine stains." In people with Parkes Weber syndrome, capillary malformations occur together with multiple micro-AVFs, which are tiny abnormal connections between arteries and veins that affect blood circulation. These AVFs can be associated with life-threatening complications including abnormal bleeding and heart failure.

Another characteristic feature of Parkes Weber syndrome is overgrowth of one limb, most commonly a leg. Abnormal growth occurs in bones and soft tissues, making one of the limbs longer and larger around than the corresponding one.

Some vascular abnormalities seen in Parkes Weber syndrome are similar to those that occur in a condition called capillary malformation-arteriovenous malformation syndrome (CM-AVM). CM-AVM and some cases of Parkes Weber syndrome have the same genetic cause.

Frequency

Parkes Weber syndrome is a rare condition; its exact prevalence is unknown.

Genetic Changes

Some cases of Parkes Weber syndrome result from mutations in the *RASA1* gene. When the condition is caused by *RASA1* gene mutations, affected individuals usually have multiple capillary malformations. People with Parkes Weber syndrome who do not have multiple capillary malformations are unlikely to have mutations in the *RASA1* gene; in these cases, the cause of the condition is often unknown.

The *RASA1* gene provides instructions for making a protein known as p120-RasGAP, which is involved in transmitting chemical signals from outside the cell to the nucleus. These signals help control several important cell functions, including the growth and division (proliferation) of cells, the process by which cells mature to carry out specific

functions (differentiation), and cell movement. The role of the p120-RasGAP protein is not fully understood, although it appears to be essential for the normal development of the vascular system.

Mutations in the *RASA1* gene lead to the production of a nonfunctional version of the p120-RasGAP protein. A loss of this protein's activity disrupts tightly regulated chemical signaling during development. However, it is unclear how these changes lead to the specific vascular abnormalities and limb overgrowth seen in people with Parkes Weber syndrome.

Inheritance Pattern

Most cases of Parkes Weber syndrome occur in people with no history of the condition in their family. These cases are described as sporadic.

When Parkes Weber syndrome is caused by mutations in the *RASA1* gene, it is sometimes inherited from an affected parent. In these cases, the condition has an autosomal dominant pattern of inheritance. Autosomal dominant inheritance means one copy of the altered gene in each cell is sufficient to cause the disorder.

Other Names for This Condition

- Parkes-Weber syndrome
- PKWS

Diagnosis & Management

Genetic Testing

 Genetic Testing Registry: Parkes Weber syndrome https://www.ncbi.nlm.nih.gov/gtr/conditions/CN074207/

Other Diagnosis and Management Resources

 GeneReview: RASA1-Related Disorders https://www.ncbi.nlm.nih.gov/books/NBK52764

General Information from MedlinePlus

- Diagnostic Tests
 https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html

- Palliative Care https://medlineplus.gov/palliativecare.html
- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html

Additional Information & Resources

MedlinePlus

- Encyclopedia: Arteriovenous Malformation Cerebral https://medlineplus.gov/ency/article/000779.htm
- Health Topic: Arteriovenous Malformations https://medlineplus.gov/arteriovenousmalformations.html
- Health Topic: Vascular Diseases https://medlineplus.gov/vasculardiseases.html

Genetic and Rare Diseases Information Center

 Parkes Weber syndrome https://rarediseases.info.nih.gov/diseases/9787/parkes-weber-syndrome

Additional NIH Resources

- National Heart, Lung, and Blood Institute: Circulation and Blood Vessels https://www.nhlbi.nih.gov/health/health-topics/topics/hhw/circulation
- National Institute of Neurological Disorders and Stroke: Arteriovenous Malformation https://www.ninds.nih.gov/Disorders/All-Disorders/Arteriovenous-Malformation-Information-Page

Educational Resources

- Boston Children's Hospital http://www.childrenshospital.org/conditions-and-treatments/conditions/parkesweber-syndrome
- Cincinnati Children's Hospital Medical Center: Arteriovenous Malformations https://www.cincinnatichildrens.org/health/a/arteriovenous
- Cincinnati Children's Hospital Medical Center: Capillary Malformations https://www.cincinnatichildrens.org/health/c/capillary
- Disease InfoSearch: Parkes Weber syndrome http://www.diseaseinfosearch.org/Parkes+Weber+syndrome/5602

- MalaCards: parkes weber syndrome http://www.malacards.org/card/parkes_weber_syndrome
- Orphanet: Parkes Weber syndrome http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=90307

Patient Support and Advocacy Resources

 National Organization for Rare Disorders (NORD): Arteriovenous Malformations https://rarediseases.org/rare-diseases/arteriovenous-malformation/

GeneReviews

 RASA1-Related Disorders https://www.ncbi.nlm.nih.gov/books/NBK52764

ClinicalTrials.gov

ClinicalTrials.gov
 https://clinicaltrials.gov/ct2/results?cond=%22Parkes+Weber+syndrome%22+OR+%22Vascular+Malformations%22

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28parkes+weber+syndrome%5BTIAB%5D%29+OR+%28pkws%5BTIAB%5D%29+OR+%28parkes-weber+syndrome%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

 PARKES WEBER SYNDROME http://omim.org/entry/608355

Sources for This Summary

- Boon LM, Mulliken JB, Vikkula M. RASA1: variable phenotype with capillary and arteriovenous malformations. Curr Opin Genet Dev. 2005 Jun;15(3):265-9. Review.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15917201
- GeneReview: RASA1-Related Disorders https://www.ncbi.nlm.nih.gov/books/NBK52764

- Lobo-Mueller E, Amaral JG, Babyn PS, Wang Q, John P. Complex combined vascular malformations and vascular malformation syndromes affecting the extremities in children. Semin Musculoskelet Radiol. 2009 Sep;13(3):255-76. doi: 10.1055/s-0029-1237692. Epub 2009 Sep 1. Review.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/19724993
- Revencu N, Boon LM, Mulliken JB, Enjolras O, Cordisco MR, Burrows PE, Clapuyt P, Hammer F, Dubois J, Baselga E, Brancati F, Carder R, Quintal JM, Dallapiccola B, Fischer G, Frieden IJ, Garzon M, Harper J, Johnson-Patel J, Labrèze C, Martorell L, Paltiel HJ, Pohl A, Prendiville J, Quere I, Siegel DH, Valente EM, Van Hagen A, Van Hest L, Vaux KK, Vicente A, Weibel L, Chitayat D, Vikkula M. Parkes Weber syndrome, vein of Galen aneurysmal malformation, and other fast-flow vascular anomalies are caused by RASA1 mutations. Hum Mutat. 2008 Jul;29(7):959-65. doi: 10.1002/humu.20746.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/18446851

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/parkes-weber-syndrome

Reviewed: August 2011 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services